

High Prevalence of *Leucocytozoon* Parasites in Nestling Northern Goshawks (*Accipiter gentilis*) in the Northern Great Basin, U.S.A.

Author(s): Michelle I. Jeffries Robert A. Miller Michelle D. Laskowski and Jay D. Carlisle

Source: Journal of Raptor Research, 49(3):294-302.

Published By: The Raptor Research Foundation

DOI: <http://dx.doi.org/10.3356/JRR-14-24.1>

URL: <http://www.bioone.org/doi/full/10.3356/JRR-14-24.1>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

HIGH PREVALENCE OF *LEUCOCYTOZOON* PARASITES IN NESTLING NORTHERN GOSHAWKS (*ACCIPITER GENTILIS*) IN THE NORTHERN GREAT BASIN, U.S.A.

MICHELLE I. JEFFRIES¹

Intermountain Bird Observatory, Department of Biological Sciences, Boise State University, 1910 University Drive, Boise, ID U.S.A. 83725

ROBERT A. MILLER

Intermountain Bird Observatory and Raptor Research Center, Department of Biological Sciences, Boise State University, 1910 University Drive, Boise, ID U.S.A. 83725

MICHELLE D. LASKOWSKI AND JAY D. CARLISLE

Intermountain Bird Observatory, Department of Biological Sciences, Boise State University, 1910 University Drive, Boise, ID U.S.A. 83725

ABSTRACT.—The Northern Goshawk (*Accipiter gentilis*) is currently listed as a sensitive species by the U.S.D.A. Forest Service. Previous research in our study area, the South Hills of the Minidoka Ranger District of the Sawtooth National Forest, Idaho, identified possible signs of parasite infections among the banded adult and nestling goshawks, which could influence their survival and breeding success. Therefore, we sought to quantify the prevalence and intensity of *Leucocytozoon* parasites among a sample of nestling goshawks in the South Hills during the 2012 breeding season. We sampled 27 nestlings from 12 nests for *Leucocytozoon* parasites by examining blood smears. All sampled nestlings were infected with *Leucocytozoon* parasites. The infection intensity ranged from 0.82–10.05 *Leucocytozoon* parasites per 1000 erythrocytes (mean \pm SE = 4.35 \pm 0.54). Using site elevation, distance-to-water, nestling age, nestling sex and nest tree species as predictor variables for infection intensity by *Leucocytozoon* parasites, we employed an information theoretic approach to select a top model to determine the presence of an effect. The top model included nest tree species as the sole predictor for infection intensity. Specifically, higher *Leucocytozoon* parasite intensity was associated with quaking aspen (*Populus tremuloides*) nest trees, as compared to lodgepole pine (*Pinus contorta*). Further research will help identify management implications for this species of concern in this high altitude forest surrounded by a shrub-steppe ecosystem.

KEY WORDS: *Northern Goshawk*; *Accipiter gentilis*; *blood parasite*, *Idaho*; *Leucocytozoon*; *parasite*.

PREVALENCIA ELEVADA DE PARÁSITOS DE *LEUCOCYTOZOON* EN POLLOS DE *ACCIPITER GENTILIS* EN LA GRAN CUENCA DEL NORTE, EEUU

RESUMEN.—Actualmente *Accipiter gentilis* está catalogada como una especie sensible por el Servicio de Bosques del Departamento de Agricultura de los Estados Unidos. Investigaciones previas en nuestro área de estudio, las South Hills del Distrito de Minidoka perteneciente al Bosque Nacional Sawtooth, Idaho, identificaron posibles signos de infecciones parasitarias entre los individuos anillados adultos y pollos de *Accipiter gentilis*, que pueden influir en su supervivencia y en el éxito reproductor. Por ello, buscamos cuantificar la prevalencia y la intensidad de la presencia de parásitos del género *Leucocytozoon* en una muestra de pollos de *A. gentilis* en las South Hills durante la época reproductiva del 2012. Analizamos muestras de sangre de 27 pollos provenientes de 12 nidos en busca de parásitos del género *Leucocytozoon*. Todos los pollos estudiados estaban infectados con parásitos *Leucocytozoon*. La intensidad de infección osciló entre 0.82–10.05 parásitos de *Leucocytozoon* por cada 1000 eritrocitos (media \pm DE = 4.35 \pm 0.54). Utilizando parámetros descriptores del lugar de cría tales como altitud, distancia al agua, edad del pollo, sexo del pollo y especie del árbol donde se ubicaba el nido como variables predictivas de la intensidad de

¹ Email address: m.jeffries011@gmail.com

infección de parásitos de *Leucocytozoon*, empleamos una aproximación basada en la teoría de la información para seleccionar el mejor modelo para determinar la presencia de algún efecto. El mejor modelo incluyó la especie del árbol como el único predictor de la intensidad de infección. Específicamente, la mayor intensidad de infección de parásitos de *Leucocytozoon* estuvo asociada con los árboles de la especie *Populus tremuloides*, comparada con *Pinus contorta*. Investigaciones posteriores contribuirán a identificar pautas de gestión para esta especie de interés en este bosque de altitud elevada rodeado por un ecosistema de estepa arbustiva.

[Traducción del equipo editorial]

The Northern Goshawk (*Accipiter gentilis*; hereafter “goshawk”) is a generalist predator occupying boreal and temperate forests of the Holarctic (Squires and Reynolds 1997). The species is currently listed as a sensitive species by the U.S.D.A. Forest Service and has been identified as a local management indicator species for the Sawtooth National Forest (D. Santini pers. comm.; U.S. Forest Service 2008, 2011). Previous research within the Sawtooth National Forest has suggested that this goshawk population may be suffering from low survival rates (Kaltenecker et al. 2004). In a 7-yr study, Kaltenecker et al. (2004) found no occurrences of young nestlings banded in the area returning to breed as adults, despite banding 74 nestlings between 1997 and 2003. Skin lesions consistent with symptoms of some parasite and fungal infections were regularly observed on both nestlings and adult females of this population during the banding process (Fig. 1; K. Hasselblad pers. comm.). Moreover, black flies (Family Simuliidae), a known vector for *Leucocytozoon* parasites, are commonly present in southern Idaho (Twin Falls County Pest Abatement District 2012) and have been frequently observed within goshawk nesting stands. Given these indicators and conditions, we determined to evaluate the prevalence and intensity of *Leucocytozoon* parasites in the Sawtooth National Forest population of Northern Goshawks.

Leucocytozoids are in the phylum Apicomplexa, a sister genera to *Haemoproteus* spp. (Outlaw and Ricklefs 2011). Both are considered a sister group to *Plasmodium* spp. (Valkiūnas 2004, Outlaw and Ricklefs 2011). *Leucocytozoon* species use black flies (Family Simuliidae) as vectors for transfer (Remple 2004). As the vector takes a blood meal from the host, the parasite is transferred from the vector’s salivary glands to the blood stream of the host. Skin lesions are created from these feeding events (Smith et al. 1998). The parasite then develops in various organs (Valkiūnas 2004). There are cases in which deaths of juvenile raptors have been attributed to severe infection and black fly lesions (Hunter

et al. 1997, Smith et al. 1998, King et al. 2010). Chronic infections of blood parasites are found in otherwise healthy, adult avian populations (Asghar et al. 2011). Survivors of infection have immune systems capable of subduing the infection to low levels, which may still produce subtle, additive effects in combination with other stresses and infections (Remple 2004, Valkiūnas 2004, Kilpatrick et al. 2006, Bensch et al. 2007). Numerous studies have attempted to relate the effects of the infection to breeding success, with conflicting results (Bensch et al. 2007, Ortego et al. 2008, Lachish et al. 2011a). Knowles et al. (2009) used a meta-regression approach on selected avian studies and found increased reproductive effort (i.e., manipulated clutch size) resulted in decreased immune function and increasing parasite levels. These findings highlight the potential cost in terms of immunosuppression with increasing reproductive output.

Prevalence and intensity of parasite infections vary due to both biotic and abiotic factors for each individual (Lachish et al. 2011b). Nestling goshawks from Germany showed increasing protozoan infection prevalence with age (Krone et al. 2005). Sex-dependent results have been conflicting in the literature (Dawson and Bortolotti 1999, Fargallo and Merino 1999, Lei et al. 2013), but Lachish et al. (2011b) found that both age and sex of Blue Tits (*Cyanistes caeruleus*) influenced the infection intensity of *Plasmodium* spp. Environmental factors also play a role in determining prevalence and intensity of blood parasites in raptors and other avian species. For example, Quillfeldt et al. (2014) suggested that nesting substrate contributed to infection status in two seabird genera. Habitat structure characteristics, such as entire tree stands, have also been related to varying parasite prevalence (Arriero et al. 2008). Zamora-Vilchis et al. (2012) found elevation to be negatively correlated with blood parasite prevalence. The distance between a nest and a water source may also be an important factor for infections, because the black fly vector requires a water source to complete its life cycle (McCreadie and



Figure 1. Photo of the throat of an adult Northern Goshawk in 2014 in the South Hills of Idaho. The photo illustrates lesions consistent with symptoms of some parasite and fungal infections.

Adler 1998). Given the potential burden that *Leucocytozoon* parasites impose on raptors, and that *Leucocytozoon* parasite prevalence varies as a function of biotic and abiotic factors, it is important to document the prevalence and intensity of these parasites when assessing the health of a raptor population.

Previous studies have documented *Leucocytozoon* parasites in goshawks in other regions and countries (Toyne and Ashford 1997, Muñoz et al. 1999, Wieliczko et al. 2003), and in one case it is thought that these parasites may have been responsible for the death of a juvenile in Maryland, U.S.A. (Ward 1975). *Leucocytozoon* parasites were previously found in a population of Sharp-shinned Hawks (*Accipiter striatus*) originating from the vicinity of the Sawtooth National Forest (Smith et al. 2004). Given this information in addition to previous reports of skin lesions, we speculated that goshawks in the area might also be at risk of infection.

Our objectives in the present study were to (1) measure *Leucocytozoon* prevalence and intensity among nestling Northern Goshawks from the population in the Sawtooth National Forest in south-central Idaho, and (2) identify potential factors that may influence the prevalence and intensity of parasitism. We hypothesized that the prevalence of

Leucocytozoon parasites in this population would be high. We further predicted that the infection intensity among nestlings may be influenced by the nest tree species, distance to water, nestling age, nestling sex, and site elevation.

METHODS

We conducted this study in the South Hills of the Sawtooth National Forest in south-central Idaho, an area that encompasses the Cassia division of the Minidoka Ranger District (41° 58.8'–42° 20'N, 113° 58.8'–114° 28.8'W, elevation: 1468–2456 masl). The Cassia division contains approximately 125 000 ha and is bordered primarily by Bureau of Land Management lands (U.S. Forest Service 2003). The naturally fragmented landscape is dominated by grasslands and mountain big sagebrush (*Artemisia tridentata vaseyana*; approximately 80%; U.S. Forest Service 1980). The remaining forested landscape consists predominantly of quaking aspen (*Populus tremuloides*), lodgepole pine (*Pinus contorta*), and subalpine fir (*Abies lasiocarpa*; U.S. Forest Service 1980).

We located goshawk nests by searching historical nesting territories and additional areas prioritized through Geographic Information System (GIS)




Figure 2. *Leucocytozoon* parasite from a blood smear of a nestling Northern Goshawk. The sample was taken in 2012 from the South Hills in south-central Idaho. The slides were fixed with absolute methanol and stained using Wright–Giemsa stain. This photo was taken using a PupilCam microscope camera through a 100× microscope lens.

analysis (Miller et al. 2013). We acquired Digital Elevation Model (DEM) data (resolution = 30 m) and stream location data from the Inside Idaho database (U.S. Geological Survey 1999, Idaho Department of Environmental Quality 2006). We used the stream data to create a 30-m-resolution raster file representing the distance of each pixel from the nearest stream location. We identified nest tree species as either lodgepole pine or aspen.

Captures and Blood Samples. To gain access to nestlings, we used climbing spurs or a climbing rope to ascend nest trees. We used a photographic key to determine nestling age (Boal 1994). We banded and collected blood samples from nestlings when they were 23–34 d old, between 28 June and 11 July of 2012. All nestlings that could be safely captured and removed from the nest were placed in a sleeve to protect their wings and lowered individually to the ground. We determined the sex of nestlings based on their tarsus size while banding (Pyle 2008) or via genetic analysis. Using a needle and syringe, we collected approximately 1 ml of blood

from the jugular vein. One drop of the blood was used to prepare a thin blood smear on individual labeled microscope slides. Two slides were created for each individual nestling to better ensure the availability of a quality blood smear sample. The rest of the blood sample was stored in Queen’s Lysis buffer for genetic analysis (Hobson et al. 1997, Brumfield and Dittmann 2012). The slides were fixed in absolute methanol and stained using Wright–Giemsa stain and buffer (Hauska et al. 1999).

Parasite Identification and Quantification. *Leucocytozoon* parasites were identified on each slide using the morphological traits described by Valkiūnas et al. (2010) with the 100× oil immersion microscope objective (Fig. 2). Once all slides were viewed and prevalence (i.e., the proportion of infected individuals to non-infected individuals) known, a Ken-A-Vision PupilCAM camera was used to record 30 – 35 unique photos of each nestling blood smear. We took photos until the viewer estimated a minimum count of 3500 erythrocytes. This is well

Table 1. Nestling age, nest elevation, and distance to water for nestling Northern Goshawks during the summer of 2012 in the South Hills of the Sawtooth National Forest, Idaho. Elevation and distance to water were calculated to the nearest 30 m. Nestling age was estimated using a photographic key.

CONTINUOUS VARIABLE	MEAN \pm SE	RANGE
Nestling age	28.4 \pm 0.6 d	23 – 34 d
Site elevation	2071 \pm 17 m	1858 – 2255 m
Distance to water	372.0 \pm 28.2 m	184.39 – 644.75 m

above previously reported erythrocyte counts of 2000 (King et al. 2010, Ortego and Cordero 2010). ImageJ software was then utilized as described by Gering and Atkinson (2004) to count the total number of erythrocytes. *Leucocytozoon* parasites in these photos were counted by consensus of two consistent viewers. We defined intensity of infection as the quantity of erythrocytes infected with *Leucocytozoon* parasites per 1000 noninfected erythrocytes. Normality was achieved by log transforming the infection intensities.

Statistical Analysis. We modeled infection intensities using a Gaussian distribution with Generalized Linear Mixed Models (GLMM; Zuur et al. 2009). We created models using various ecologically relevant combinations of fixed effects (nestling sex, nestling age, site elevation, distance-to-water, and nest tree species) and a random effect of nest ID to predict the log-transformed intensity of parasite infection. We ranked models using the Akaike Information Criterion adjusted for small sample size (AIC_c ; Burnham and Anderson 2002). We measured the presence of an effect if the coefficient for a given predictor was represented in the top model and the top model was ranked above the null model (Burnham and Anderson 2002).

We conducted all statistical analyses in R (R Development Core Team 2012). We fit GLMM models using the R package “lme4” (Bolker et al. 2013). We calculated AIC_c and related values using the R package “AICcmodavg” (Mazerolle 2012). Standard error was calculated and reported as \pm from the mean.

This study was conducted with all necessary permits and authorizations including a sub-permit (RAM) under master Federal Bird Banding Permit #21633, a State of Idaho Wildlife Collection/Banding/Possession permit #870115, with authorization from Idaho Fish and Game Jerome Field Of-

Table 2. Nestling sex and nest tree species for nestling Northern Goshawks during the summer of 2012 in the South Hills of the Sawtooth National Forest, Idaho. Sex was determined through band size and genetic analysis.

NEST TREE SPECIES	SEX	
	MALE	FEMALE
Aspen	7	9
Lodgepole pine	6	5

fice and under Boise State University Institutional Animal Care and Use Committee permit #006-AC11-004.

RESULTS

We sampled 27 nestlings from 12 separate nests. Only one of these nests was occupied in the previous year. We measured the continuous variables site elevation, distance-to-water, and nestling age for each nest (Table 1). We categorized samples by sex and nest tree species (Table 2). All 27 sampled nestlings were positive for *Leucocytozoon* parasites. The intensity varied from 0.82 to 10.05 *Leucocytozoon* parasites per 1000 erythrocytes, with a mean of 4.35 ± 0.54 . We also observed unconfirmed infections of *Haemoproteus* spp. or *Plasmodium* spp. in the blood of three nestlings.

Our top model included nest tree species as the sole predictor for infection intensity (Table 3). Specifically, we found that increased *Leucocytozoon* parasite intensity was associated with aspen nesting substrate (Fig. 3).

Table 3. Generalized Linear Mixed Models used to predict *Leucocytozoon* parasite infection with a suite of environmental parameters. These parameters included fixed effects of nestling age, nestling sex, nest tree species, site elevation, and distance to water. We also included a random effect of nest ID. We ranked models using the Akaike Information Criterion adjusted for small sample size (AIC_c). The top model included nest tree species as the sole predictor for infection intensity.

MODEL	K	AIC_c	ΔAIC_c	w_i
Nest tree species	4	57.35	0.00	0.44
Null	3	58.55	1.20	0.24
Distance-to-water	4	60.31	2.96	0.10
Nestling age	4	60.88	3.52	0.08
Nestling sex	4	61.03	3.68	0.07
Site elevation	4	61.26	3.91	0.06




Figure 3. Model results indicating that increased *Leucocytozoon* parasite intensity was associated with aspen nest tree species as compared with lodgepole pine substrate. The point indicates the mean. The bold line indicates the median. The box shows the quartiles. The whiskers designate the range.

DISCUSSION

All nestling goshawks sampled in our study were infected with *Leucocytozoon* parasites. Although similarly high prevalence values for *Leucocytozoon* parasites have been reported in nestlings of other raptor species (90%, 100%; Forrester et al. 2001, King et al. 2010), prior studies on nestling goshawks have reported much lower values (18.8%, 53.6%; Toyne and Ashford 1997, Wieliczko et al. 2003). Due to the dearth of research showing a high prevalence of *Leucocytozoon* parasites in goshawks, this population is of interest.

Although the goshawks in our study showed a high prevalence of *Leucocytozoon* parasites among the population of nestlings, it was unclear to what degree these parasites may have negatively affected the population. In 2011 and 2012, occupancy rate and reproductive rate in the study area were equal to or above average (Bechard et al. 2006, Kenward 2006, Miller et al. 2014), and we now have evidence that nestlings fledged in the South Hills have survived to reproduce in the area. These cases include three adult female breeders that hatched and later bred in this area. The observed subadult breeding rate in the area was <10%, which further suggested the population was stable (R. Miller unpubl. data, Kenward 2006).

The infection intensity varied from 0.82 to 10.05 *Leucocytozoon* parasites per 1000 erythrocytes, with a mean of 4.35 ± 0.54 . This intensity range and

average was lower than that in other studies of nestling raptors (1–87.5, 0–75; King et al. 2010, Ortego and Cordero 2010). This may indicate that this population managed infections to lower levels. There are a number of possible explanations for how this population of goshawks may manage the effects of infection by *Leucocytozoon* parasites. First, food availability may be sufficiently high that nestlings are able to overcome the debilitating effects of parasites. In a study of Great Horned Owls (*Bubo virginianus*), Hunter et al. (1997) found that prey shortages greatly increased the effect of infection by *Leucocytozoon* parasites on fledgling survival. In years of lower than average food supply, birds may allocate more energy to maintaining body condition, rather than to immune defenses, leaving them more susceptible to parasites. Although Miller et al. (2014) found that prey consumption per nestling goshawk in our study area in 2011 and 2012 was similar to or greater than that in other studies (Younk and Bechard 1994, Smithers et al. 2005), the limited duration of our study precluded a test of the effects of different levels of prey abundance.

The stability of a goshawk population in the face of high *Leucocytozoon* parasite prevalence may also be a consequence of nest preference and supplementation behaviors by breeding individuals. Kenward (2006) suggested that intermittent use of nesting structures decreases the buildup of parasites. Having fewer parasitic stressors indirectly allows nestlings to better respond to *Leucocytozoon* parasite infections. Of the 12 nests we sampled in 2012, only one was known to be occupied in 2011. This nest also fell within the upper 74th percentile of infection intensity, consistent with Kenward's hypothesis. We also observed that adult goshawks delivered fresh aspen and lodgepole pine branches to add to their nests multiple times each day. These branches buried uneaten prey items and unhatched goshawk eggs (R. Miller unpub. data), which reduced exposure of decaying waste and perhaps attraction of black flies. Resin from the broken sprigs may have repelled the black flies due to the presence of volatile chemicals (Lafuma et al. 2001, Keeling and Bohlmann 2006, Lindroth and Clair 2013). Although all nestlings in our study were infected by *Leucocytozoon* parasites, any reduction in fly activity and harassment may lessen the negative effects of these flies (Smith et al. 1998).

Our top model included nest tree species as the sole predictor for infection intensity. Specifically, higher *Leucocytozoon* parasite intensity was associated

with aspen nesting substrate rather than with lodgepole pine substrate. Both conifers and aspen contain volatile compounds, and perhaps conifers are more efficient at deterring black flies (Lafuma et al. 2001, Keeling and Bohlmann 2006, Lindroth and Clair 2013). Lodgepole pine branches were the species most frequently deposited in the nests, even for nests located in aspen trees (R. Miller unpubl. data), which supported this idea. Aspen trees are suitable nesting substrate for these goshawks and may provide other advantages, but may be associated with a cost: higher *Leucocytozoon* parasite infection intensities.

A number of factors may have contributed to our inability to detect strong relationships between *Leucocytozoon* parasite infection intensity and the various other variables we measured. First, there may have been insufficient variation in infection intensities to detect effects across the range of variables we measured using our available sample size. For example, the nearly ubiquitous access to water, brief appropriate sampling age of nestlings, and the relatively narrow elevation band suitable for nesting at our study site may not have generated enough variation in these parameters to manifest substantial changes in infection intensity. Selecting nests that fall within a greater range of these parameters, assuming goshawks nest in these areas, may improve the resolution of the model. Increasing sample size may also help to address problems associated with low power. Lastly, examining how vector (i.e., Family Simuliidae) distribution, rather than surrogate measures such as elevation and distance-to-water, relates to infection intensity might provide greater insight into variation in infection intensity.

Our results demonstrate that there is a high prevalence of *Leucocytozoon* parasites in this population of goshawks, and that the birds seem to be enduring nonetheless. The low infection intensity range and rate may indicate that *Leucocytozoon* parasites are not a limitation to this population. However, further study is needed to more thoroughly understand the consequences of parasitism in this host-parasite relationship. For example, are the effects of parasitism more prominent in years of food and climate stress? What environmental variables explain variation in vector abundance? Unraveling the details of the relationship between *Leucocytozoon* parasites and goshawks will help address management concerns for this species, as well as other avian species living in this high altitude forest surrounded by a shrub-steppe ecosystem.

ACKNOWLEDGMENTS

Special thanks to U.S.D.A. Forest Service, Minidoka Ranger District, particularly wildlife biologist Dena Santini, for their financial, consulting, and equipment support for the completion of this project. Additional funding was received from a Michael W. Butler Ecological Research Award to R. Miller. Logistic and equipment support was received from Boise State University's Raptor Research Center, the Intermountain Bird Observatory, the Department of Biological Sciences, and David L. Anderson. We received statistical consultation from Boise State University statistician Laura Bond. We also thank Ian Robertson for his review of this manuscript. Much of this work was completed using volunteer time from field volunteers including: David L. Anderson, Alexis Billings, Karyn deKramer, Greg Kaltenecker, Kraig Laskowski, Thurman Pratt, Emmy Tyrrell, Heidi Ware, and Mike Zinn.

LITERATURE CITED

- ARRIERO, E., J. MORENO, S. MERINO, AND J. MARTÍNEZ. 2008. Habitat effects on physiological stress response in nestling Blue Tits are mediated through parasitism. *Physiological and Biochemical Zoology* 81:195–203.
- ASGHAR, M., D. HASSELQUIST, AND S. BENSCH. 2011. Are chronic avian haemosporidian infections costly in wild birds? *Journal of Avian Biology* 42:530–537.
- BECHARD, M.J., G.D. FAIRHURST, AND G.S. KALTENECKER. 2006. Occupancy, productivity, turnover and dispersal of Northern Goshawks in portions of the northeastern Great Basin. *Studies in Avian Biology* 31:100–108.
- BENSCH, S., J. WALDENSTRÖM, N. JONZÉN, H. WESTERDAHL, B. HANSSON, D. SEJBERG, AND D. HASSELQUIST. 2007. Temporal dynamics and diversity of avian malaria parasites in a single host species. *Journal of Animal Ecology* 76:112–122.
- BOAL, C.W. 1994. A photographic and behavioral guide to aging nestling Northern Goshawks. *Studies in Avian Biology* 16:32–40.
- BOLKER, B., D. BATES, M. MAECHLER, AND S. WALKER. 2013. lme4: linear mixed-effects models using Eigen and S4 (version 1.0-6). <http://lme4.r-forge.r-project.org/> (last accessed 9 May 2014).
- BRUMFIELD, R. AND D. DITTMANN. 2012. Recipes for DNA Buffers. http://aou.org/committees/collections/recipes_dna_buffers.php (last accessed 17 December 2012).
- BURNHAM, K. AND D. ANDERSON. 2002. Model selection and multi-model inference: a practical information-theoretic approach. Springer-Verlag, New York, NY U.S.A.
- DAWSON, R.D. AND G.R. BORTOLOTTI. 1999. Prevalence and intensity of hematozoan infections in a population of American Kestrels. *Canadian Journal of Zoology* 77:162–170.
- FARGALLO, J.A. AND S. MERINO. 1999. Brood size manipulation modifies the intensity of infection by Haematozoa in female Blue Tits *Parus caeruleus*. *Ardea* 87:261–268.
- FORRESTER, D.J., G.W. FOSTER, AND J.L. MORRISON. 2001. *Leucocytozoon toddi* and *Haemoproteus tinnunculi* (Protozoa: Haemosporina) in the Chimango Caracara (*Milvago chimango*) in southern Chile. *Memórias do Instituto Oswaldo Cruz* 96:1023–1024.

- GERING, E. AND C.T. ATKINSON. 2004. A rapid method for counting nucleated erythrocytes on stained blood smears by digital image analysis. *Journal of Parasitology* 90:879–881.
- HAUSKA, H., A. SCOPE, L. VASICEK, AND B. REAUZ. 1999. Comparison of different staining methods for avian blood cells. *Tierärztliche Praxis. Ausgabe K, Kleintiere/Heimtiere* 27:280–287.
- HOBSON, K.A., H.L. GIBBS, AND M.L. GLOUTNEY. 1997. Preservation of blood and tissue samples for stable-carbon and stable-nitrogen isotope analysis. *Canadian Journal of Zoology* 75:1720–1723.
- HUNTER, D.B., C. ROHNER, AND D.C. CURRIE. 1997. Mortality in fledgling Great Horned Owls from black fly hematophaga and leucocytozoonosis. *Journal of Wildlife Diseases* 33:486–491.
- IDAHO DEPARTMENT OF ENVIRONMENTAL QUALITY. 2006. Streams of Idaho (2002 305(b) and 303(d) integrated report—water quality). Idaho Department of Environmental Quality, Boise, ID U.S.A. http://cloud.insideidaho.org/data/ideq/strm305b2002_id_ideq (last accessed 28 October 2012).
- KALTENECKER, G.S., K.W. HASSELBLAD, M.J. BECHARD, AND J.W. BEALS. 2004. Occupancy, productivity, and banding of Northern Goshawks on the Sawtooth National Forest. 2003. Idaho Department of Fish and Game, Boise, ID U.S.A.
- KEELING, C.I. AND J. BOHLMANN. 2006. Genes, enzymes and chemicals of terpenoid diversity in the constitutive and induced defence of conifers against insects and pathogens. *New Phytologist* 170:657–675.
- KENWARD, R. 2006. The goshawk. T. and A.D. Poyser, London, U.K.
- KILPATRICK, A.M., D.A. LAPOINTE, C.T. ATKINSON, B.L. WOODWORTH, J.K. LEASE, M.E. REITER, AND K. GROSS. 2006. Effects of chronic avian malaria (*Plasmodium relictum*) infection on reproductive success of Hawaii Amakihi (*Hemignathus virens*). *Auk* 123:764–774.
- KING, J.C., S.A. DUBAY, T.C. HUSPENI, A.R. VANLANEN, AND R.W. GERHOLD. 2010. Parasite infections in nestling Red-shouldered Hawks (*Buteo lineatus*) in northeast Wisconsin. *Journal of Parasitology* 96:535–540.
- KNOWLES, S.C.L., S. NAKAGAWA, AND B.C. SHELDON. 2009. Elevated reproductive effort increases blood parasitaemia and decreases immune function in birds: a meta-regression approach. *Functional Ecology* 23:405–415.
- KRONE, O., R. ALTENKAMP, AND N. KENNTNER. 2005. Prevalence of *Trichomonas gallinae* in Northern Goshawks from the Berlin area of northeastern Germany. *Journal of Wildlife Disease* 41:304–309.
- LACHISH, S., S.C.L. KNOWLES, R. ALVES, M.J. WOOD, AND B.C. SHELDON. 2011a. Infection dynamics of endemic malaria in a wild bird population: parasite species-dependent drivers of spatial and temporal variation in transmission rates. *Journal of Animal Ecology* 80:1207–1216.
- , ———, ———, ———, AND ———. 2011b. Fitness effects of endemic malaria infections in a wild bird population: the importance of ecological structure. *Journal of Animal Ecology* 80:1196–1206.
- LAFUMA, L., M.M. LAMBRECHTS, AND M. RAYMOND. 2001. Aromatic plants in bird nests as protection against blood-sucking flying insects? *Behavioural Processes* 56:113–120.
- LEI, B., A. AMAR, A. KOESLAG, T.A. GOUS, AND G.J. TATE. 2013. Differential hemoparasite intensity between Black Sparrowhawk (*Accipiter melanoleucus*) morphs suggests an adaptive function for polymorphism. *PLoS ONE* 8:e81607.
- LINDROTH, R.L. AND S.B. ST. CLAIR. 2013. Adaptations of quaking aspen (*Populus tremuloides* Michx.) for defense against herbivores. *Forest Ecology and Management* 299:14–21.
- MAZEROLLE, M.J. 2012. AICmodavg: model selection and multimodel inference based on (Q)AIC(c). Version 1.26. <http://CRAN.R-project.org/package=AICmodavg> (last accessed 9 May 2014).
- MCCREADIE, J.W. AND P.H. ADLER. 1998. Scale, time, space, and predictability: species distributions of preimaginal black flies (Diptera: Simuliidae). *Oecologia* 114:79–92.
- MILLER, R.A., J.D. CARLISLE, AND M.J. BECHARD. 2014. Effects of prey abundance on breeding season diet of Northern Goshawks (*Accipiter gentilis*) within an unusual prey landscape. *Journal of Raptor Research* 48:1–12.
- , ———, ———, AND D. SANTINI. 2013. Predicting nesting habitat of Northern Goshawks in mixed aspen-lodgepole pine forests in a high-elevation shrub-steppe dominated landscape. *Open Journal of Ecology* 3:109–115.
- MUÑOZ, E., D. FERRER, R. MOLINA, AND R.D. ADLARD. 1999. Prevalence of haematozoa in birds of prey in Catalonia, north-east Spain. *Veterinary Record* 144:632–636.
- ORTEGO, J. AND P.J. CORDERO. 2010. Factors associated with the geographic distribution of leucocytozoa parasitizing nestling Eagle Owls (*Bubo bubo*): a local spatial-scale analysis. *Conservation Genetics* 11:1479–1487.
- , ———, J.M. APARICIO, AND G. CALABUIG. 2008. Consequences of chronic infections with three different avian malaria lineages on reproductive performance of Lesser Kestrels (*Falco naumanni*). *Journal of Ornithology* 149:337–343.
- OUTLAW, D.C. AND R.E. RICKLEFS. 2011. Rerooting the evolutionary tree of malaria parasites. *Proceedings of the National Academy of Sciences of the United States of America* 108:13183–13187.
- PYLE, P. 2008. Identification guide to North American birds. Part 2. Slate Creek Press, Bolinas, CA U.S.A.
- QUILLFELDT, P., J. MARTÍNEZ, L. BUGONI, P.L. MANCINI, AND S. MERINO. 2014. Blood parasites in noddies and boobies from Brazilian offshore islands—differences between species and influence of nesting habitat. *Parasitology* 141:399–410.

- R DEVELOPMENT CORE TEAM. 2012. R: a language and environment for statistical computing. Vienna, Austria. <http://www.R-project.org> (last accessed 10 December 2012).
- REMPEL, J.D. 2004. Intracellular hematozoa of raptors: a review and update. *Journal of Avian Medicine and Surgery* 18:75–88.
- SMITH, R.B., E.C. GREINER, AND B.O. WOLF. 2004. Migratory movements of Sharp-shinned Hawks (*Accipiter striatus*) captured in New Mexico in relation to prevalence, intensity, and biogeography of avian hematozoa. *Auk* 121:837–846.
- SMITH, R.N., S.L. CAIN, S.H. ANDERSON, J.R. DUNK, AND E.S. WILLIAMS. 1998. Blackfly-induced mortality of nestling Red-tailed Hawks. *Auk* 115:368–375.
- SMITHERS, B.L., C.W. BOAL, AND D.E. ANDERSEN. 2005. Northern Goshawk diet in Minnesota: an analysis using video recording systems. *Journal of Raptor Research* 39:264–273.
- SQUIRES, J.R. AND R.T. REYNOLDS. 1997. Northern Goshawk (*Accipiter gentilis*). In A. Poole [Ed.], *The birds of North America* online, No. 298. Cornell Lab of Ornithology, Ithaca, NY U.S.A. <http://bna.birds.cornell.edu/bna/species/298> (last accessed 20 October 2012).
- TOYNE, E.P. AND R.W. ASHFORD. 1997. Blood parasites of nestling goshawks. *Journal of Raptor Research* 31:81–83.
- TWIN FALLS COUNTY PEST ABATEMENT DISTRICT 2012. Why all the black flies? Twin Falls, ID U.S.A. <http://www.tfcpad.qwestoffice.net/> (last accessed 20 January 2014).
- U.S. FOREST SERVICE. 1980. Cassia timber environmental assessment. Sawtooth National Forest, Twin Falls, ID U.S.A.
- . 2003. Sawtooth National Forest revised land and resource management plan. Twin Falls, ID U.S.A.
- . 2008. Intermountain region proposed, endangered, threatened, and sensitive species. Ogden, UT U.S.A. http://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5370041.pdf (last accessed 1 April 2015).
- . 2011. Sawtooth National Forest terrestrial management indicator species forest plan 5-year monitoring report (2011) for Pileated Woodpecker and Greater Sage-Grouse. Twin Falls, ID U.S.A.
- U.S. GEOLOGICAL SURVEY. 1999. National elevation dataset of southcentral Idaho. U.S. Geological Survey, Sioux Falls, SD U.S.A. http://cloud.insideidaho.org/data/elevation/ned/1999_30m_idaho/ned30m_sc (last accessed 17 November 2010).
- VALKIUNAS, G. 2004. Avian malaria parasites and other haemosporidia. CRC Press, Boca Raton, FL U.S.A.
- , R.N.M. SEHGAL, T.A. IEZHOVA, AND A.C. HULL. 2010. Identification of *Leucocytozoon toddi* group (Haemosporida: Leucocytozoidae), with remarks on the species taxonomy of leucocytozoids. *Journal of Parasitology* 96:170–177.
- WARD, F.P. 1975. A clinical evaluation of parasites of birds of prey. *Journal of Zoo Animal Medicine* 6:3–8.
- WIELICZKO, A., T. PIASECKI, G.M. DORRESTEIN, A. ADAMSKI, AND M. MAZURKIEWICZ. 2003. Evaluation of the health status of goshawk chicks (*Accipiter gentilis*) nesting in Wrocław vicinity. *Bulletin of the Veterinary Institute in Pulawy* 47:247–257.
- YOUNK, J.V. AND M.J. BECHARD. 1994. Breeding ecology of the Northern Goshawk in high-elevation aspen forests of northern Nevada. *Studies in Avian Biology* 16:119–121.
- ZAMORA-VILCHIS, I., S.E. WILLIAMS, AND C.N. JOHNSON. 2012. Environmental temperature affects prevalence of blood parasites of birds on an elevation gradient: implications for disease in a warming climate. *PLoS ONE* 7:e39208.
- ZUUR, A.F., E.N. IENO, N.J. WALKER, A.A. SAVELIEV, AND G.M. SMITH. 2009. *Mixed effects models and extensions in ecology with R*. Springer, New York, NY U.S.A.

Received 12 March 2014; accepted 16 December 2014

Associate Editor: Pascual López-López